

Submitted to MNRAS

The cosmic rate of Pair-Instability Supernovae

Francesco Gabrielli, Andrea Lapi, Lumen Boco, Cristiano Ugolini, Guglielmo Costa, Cecilia Sgalletta, Kendall Shepherd, Ugo Niccolò Di Carlo, Alessandro Bressan, Marco Limongi, Mario Spera




very-massive stars

low metallicity

$Z < Z_{\odot}/2$ Langer et al. 2007

$Z < Z_{\odot}/3$ Higgins et al. 2021



$100 M_{\odot}$

$140 M_{\odot}$

$260 M_{\odot}$

$300 M_{\odot}$

very-massive stars

low metallicity

$Z < Z_{\odot}/2$ Langer et al. 2007

$Z < Z_{\odot}/3$ Higgins et al. 2021



$100 M_{\odot}$

$140 M_{\odot}$

$260 M_{\odot}$


$300 M_{\odot}$

very-massive stars

low metallicity

$Z < Z_{\odot}/2$ Langer et al. 2007

$Z < Z_{\odot}/3$ Higgins et al. 2021



$100 M_{\odot}$

$140 M_{\odot}$

$260 M_{\odot}$


$300 M_{\odot}$

very-massive stars

low metallicity

$Z < Z_{\odot}/2$ Langer et al. 2007

$Z < Z_{\odot}/3$ Higgins et al. 2021



PPISN

$100 M_{\odot}$

$140 M_{\odot}$

$260 M_{\odot}$


$300 M_{\odot}$

very-massive stars

low metallicity

$Z < Z_{\odot}/2$ Langer et al. 2007

$Z < Z_{\odot}/3$ Higgins et al. 2021



PPISN

PISN

$100 M_{\odot}$

$140 M_{\odot}$

$260 M_{\odot}$


$300 M_{\odot}$

very-massive stars

low metallicity

$Z < Z_{\odot}/2$ Langer et al. 2007

$Z < Z_{\odot}/3$ Higgins et al. 2021



PPISN

PISN

DCBH

$100 M_{\odot}$

$140 M_{\odot}$

$260 M_{\odot}$

$300 M_{\odot}$

no PISN observations

only candidate identifications
(e.g. Super-Luminous Supernovae, SLSNe)

no PISN observations

only candidate identifications
(e.g. Super-Luminous Supernovae, SLSNe)

**several hundreds
CCSN observations**

$$L_{\text{PISN}} \lesssim 10^2 L_{\text{CCSN}}$$

no PISN observations

only candidate identifications
(e.g. Super-Luminous Supernovae, SLSNe)

**several hundreds
CCSN observations**

$$L_{\text{PISN}} \lesssim 10^2 L_{\text{CCSN}}$$

where are the PISNe?

host galaxies?

no PISN observations

only candidate identifications
(e.g. Super-Luminous Supernovae, SLSNe)

**several hundreds
CCSN observations**

$$L_{\text{PISN}} \lesssim 10^2 L_{\text{CCSN}}$$

where are the PISNe?

host galaxies?

intrinsically few

observational issues

no PISN observations

only candidate identifications
(e.g. Super-Luminous Supernovae, SLSNe)

**several hundreds
CCSN observations**

$$L_{\text{PISN}} \lesssim 10^2 L_{\text{CCSN}}$$

where are the PISNe?

host galaxies?

intrinsically few

observational issues

Aim

cosmic PISN rate density

$$\frac{dN_{PISN}}{dt dV}(z) = \int dZ \frac{dM_{SFR}}{dt dV dZ}(z, Z) \times \frac{dN_{PISN}}{dM_{SFR}}(Z)$$

Aim

cosmic PISN rate density

$$\frac{dN_{PISN}}{dt dV}(z) = \int dZ \frac{dM_{SFR}}{dt dV dZ}(z, Z) \times \frac{dN_{PISN}}{dM_{SFR}}(Z)$$

explore
dependence on

galaxy evolution model

stellar evolution
simulations

Methods

Galaxy semi-empirical model

Galaxy semi-empirical model


**Z-dependent
Star Formation Rate Density
(SFRD)**

$$\frac{dM_{SFR}(z, Z)}{dt dV dZ}$$

Galaxy semi-empirical model

Z-dependent
Star Formation Rate Density
(SFRD)

$$\frac{dM_{SFR}(z, Z)}{dt dV dZ}$$



Galaxy semi-empirical model


Z-dependent
Star Formation Rate Density
(SFRD)

$$\frac{dM_{SFR}(z, Z)}{dt dV dZ}$$

$$\sigma_Z = 0.15$$

$$0.35$$

$$0.70$$



Stellar evolution model

$$\frac{dN_{PISN}}{dM_{SFR}}(Z) \propto \int_{M_{entry}}^{M_{exit}} \phi(M) dM$$

Stellar evolution model

$$\frac{dN_{PISN}}{dM_{SFR}}(Z) \propto \int_{M_{entry}}^{M_{exit}} \phi(M) dM$$



Kroupa IMF

$[0.1 M_{\odot}, M_{up}]$

$150 M_{\odot}$

$300 M_{\odot}$

Stellar evolution model

$$\frac{dN_{PISN}}{dM_{SFR}}(Z) \propto \int_{M_{entry}}^{M_{exit}} \phi(M) dM$$

Kroupa IMF

$[0.1 M_{\odot}, M_{up}]$

$150 M_{\odot}$
 $300 M_{\odot}$

$[45, 120] M_{\odot}$
 $[55, 110] M_{\odot}$
 $[60, 105] M_{\odot}$

M_{CO}

$M_{entry/exit}(Z)$

PARSEC

FRANEC

Bressan et al. 2012, Costa et al. 2019, 2021
Spera & Mapelli 2017, Iorio et al 2022

Chieffi & Limongi 2013, Limongi & Chieffi 2018

Stellar evolution model

single stars

$$\frac{dN_{PISN}}{dM_{SFR}}(Z) \propto \int_{M_{entry}}^{M_{exit}} \phi(M) dM$$

Kroupa IMF

$$[0.1 M_{\odot}, M_{up}]$$

150 M_{\odot}

300 M_{\odot}

[45, 120] M_{\odot}

[55, 110] M_{\odot}

[60, 105] M_{\odot}

M_{CO}

$M_{entry/exit}(Z)$

PARSEC

FRANEC


Bressan et al. 2012, Costa et al. 2019, 2021
Spera & Mapelli 2017, Iorio et al 2023

Chieffi & Limongi 2013, Limongi & Chieffi 2018

single stars

$$\frac{dN_{PISN}}{dM_{SFR}}(Z)$$

stellar variations




name	stellar code	M_{CO}/M_{\odot}	M_{up}/M_{\odot}	Z_{max}
P	FRANEC	60-105	150	1.5×10^{-3}
M1	PARSEC-I	55-110	150	1.5×10^{-3}
M2	FRANEC	45-120	150	5.5×10^{-3}
F	PARSEC-I	55-110	300	6.6×10^{-3}
M3	PARSEC-II	45-120	150	1.0×10^{-2}
O	PARSEC-II	45-120	300	1.7×10^{-2}

single stars

$$\frac{dN_{PISN}}{dM_{SFR}}(Z)$$

stellar variations



name	stellar code	M_{CO}/M_{\odot}	M_{up}/M_{\odot}	Z_{max}
P	FRANEC	60-105	150	1.5×10^{-3}
M1	PARSEC-I	55-110	150	1.5×10^{-3}
M2	FRANEC	45-120	150	5.5×10^{-3}
F	PARSEC-I	55-110	300	6.6×10^{-3}
M3	PARSEC-II	45-120	150	1.0×10^{-2}
O	PARSEC-II	45-120	300	1.7×10^{-2}

PISNe in binaries

PARSEC
(Bressan et al. 2012, Costa et al. 2019, 2021)

SEVN
(Iorio et al 2023)



population of
single stars

population of
binaries

$$f_{bin} = 0.50$$

PISNe in binaries


PARSEC
(Bressan et al. 2012, Costa et al. 2019, 2021)

SEVN
(Iorio et al 2023)

population of
single stars

population of
binaries


$$f_{bin} = 0.50$$



$$f_{bin}^{PISN} = 0.56$$

Results

stellar variations




name	stellar code	M_{CO}/M_{\odot}	M_{up}/M_{\odot}	Z_{max}
P	FRANEC	60-105	150	1.5×10^{-3}
M1	PARSEC-I	55-110	150	1.5×10^{-3}
M2	FRANEC	45-120	150	5.5×10^{-3}
F	PARSEC-I	55-110	300	6.6×10^{-3}
M3	PARSEC-II	45-120	150	1.0×10^{-2}
O	PARSEC-II	45-120	300	1.7×10^{-2}

galactic variations

Z_{max}



σ_Z




σ_Z



$\sigma_Z = 0.15$

$\sigma_Z = 0.35$


$\sigma_Z = 0.70$



Z_{max}




σ_Z




Z_{max}

σ_Z




Z_{max}



host galaxy properties


variation F
 $\sigma_Z = 0.15$



host galaxy properties

$$M_{\star} \sim 10^9 - 10^{10} M_{\odot}$$


$$\text{variation } F$$
$$\sigma_Z = 0.15$$



host galaxy properties



$$M_{\star} \sim 10^9 - 10^{10} M_{\odot}$$

$$\text{variation } F \\ \sigma_Z = 0.15$$



$$Z_{max} = 8 \times 10^{-3}$$

$$Z \lesssim Z_{max}$$



peak of PISN rate at
 $Z \sim 10^{-3} - 10^{-2}$



Pop II stars main PISN progenitors, not Pop III

CAVEAT: SFRD and IMF at high z highly uncertain

Conclusions

strong dependence on
stellar and galactic variations

→ PISN rate down to $\sim 10^{-4}/\text{yr Gpc}^3$ ($z = 0$)

↓
intrinsically few

Conclusions

strong dependence on
stellar and galactic variations

→ PISN rate down to $\sim 10^{-4}/\text{yr Gpc}^3$ ($z = 0$)

↓
intrinsically few

possible (or lack of) future PISN observations could pose constraints on

→ maximum stellar Z to have PISN
upper limit of stellar IMF
dispersion of galaxy Z distribution in z

PISN contribution from binaries similar to single stars

PISN contribution from binaries similar to single stars


main contribution to PISN rate from $Z \sim 10^{-3} - 10^{-2}$



Pop II stars main PISN progenitors, not Pop III

PISN contribution from binaries similar to single stars

main contribution to PISN rate from $Z \sim 10^{-3} - 10^{-2}$



Pop II stars main PISN progenitors, not Pop III

single galaxy (M_{\star} , Z , ψ , z) contribution to PISN rate



possible indications to future observational campaigns from host galaxy properties

Follow-up

where are the PISNe?

```
graph TD; A[where are the PISNe?] --> B[intrinsically few]; A --> C[observational issues]; C --- D[PISN detection rate with JWST];
```

intrinsically few

observational issues

PISN detection rate with JWST

Back-up slides

Possible reasons for missed observation

assuming theory of stellar evolution is correct

PISNe only at high z (low Z environments) \longrightarrow too dim to be observed

stellar Initial Mass Function (IMF) does not extend up to PISN range
stars with mass $> 300 M_{\odot}$ observed (stellar mergers?)

PISNe preferentially in dusty environments \longrightarrow emission blocked by dust

PISNe more rare than CCSNe

PISN emission

early times

conversion of kinetic and radiation energy
into thermal energy

later times

large amounts of radioactive ^{56}Ni
up to $\sim 60 M_{\odot}$

near-infrared band


expected PISN luminosity

$$L_{PISN} \lesssim 10^{44} \text{ erg/s}$$

typical luminosity of
Core-Collapse Supernovae (CCSNe)
routinely observed in Local Universe


$$L_{CCSN} \sim 10^{42} \text{ erg/s}$$

Galaxy semi-empirical model




Galaxy semi-empirical model

Galaxy Stellar Mass Functions (GSMF)




Davidson et al. 2017, Weaver et al. 2023

Galaxy Main Sequence (MS)



Speagle et al. 2014, Popesso et al. 2023

Fundamental Metallicity Relation (FMR)



Mannucci et al. 2010, Curti et al. 2020, 2023

Galaxy Stellar Mass Functions

Chruslinska and Nelemans 2019




analytical fits to observations

$$\Phi(M_{\star}) \propto e^{-M_{\star}/M_c} \left(\frac{M_{\star}}{M_c} \right)^{\alpha_{GSMF}}$$

combine several determinations

low-mass end slope


$$\begin{aligned} \alpha_{GSMF} &= -1.45 \\ &= -0.1z - 1.34 \end{aligned}$$



Galaxy Main Sequence


$$\frac{dp}{d \log \psi}(\psi, M_{\star}, z) = \frac{f_{MS}}{\sigma_{MS} \sqrt{2\pi}} \exp \left[-\frac{(\log \psi - \langle \log \psi \rangle_{MS})^2}{2\sigma_{MS}^2} \right] + \frac{f_{SB}}{\sigma_{SB} \sqrt{2\pi}} \exp \left[-\frac{(\log \psi - \langle \log \psi \rangle_{SB})^2}{2\sigma_{SB}^2} \right]$$

Popesso et al. 2022



$$\log \psi = (-27.58 + 0.26t) + (4.95 - 0.04t) \log M_{\star} - 0.2(\log M_{\star})^2$$


Chruslinska et al. 2021



$$\begin{aligned} f_{MS} + f_{SB} &= 1 \\ \langle \log \psi \rangle_{SB} &= \langle \log \psi \rangle_{MS} + 0.59 \\ \sigma_{MS} &= 0.188 \quad \sigma_{SB} = 0.243 \end{aligned}$$

Z evolution

$$Z_{FMR}(M_{\star}, \psi)$$



Curti et al. 2020

$$Z(M, \text{SFR}) = Z_0 - \gamma/\beta \log(1 + (M/M_0(\text{SFR}))^{-\beta})$$

$$\log(M_0(\text{SFR})) = m_0 + m_1 \log(\text{SFR})$$

Z_0	m_0	m_1	γ	β
8.779 ± 0.005	10.11 ± 0.03	0.56 ± 0.01	0.31 ± 0.01	2.1 ± 0.4

$$\log Z = 12 + \log(\text{O}/\text{H}) - 10.58$$


$$\frac{dp}{d \log Z}(Z, Z_{FMR}) \propto \exp \left[-\frac{(\log Z - \log Z_{FMR})^2}{2\sigma_Z^2} \right]$$

$$\sigma_Z = [0.15, 0.35, 0.70]$$

$M_{\text{entry/exit}}$ ranges

$Z \backslash M_{\text{CO}}$	1×10^{-4}	1×10^{-3}	4×10^{-3}	8×10^{-3}	1×10^{-2}	2×10^{-2}
PARSEC-I						
45-120	108-257	109-435	158-458	178-222	-	-
55-110	126-237	128-382	195-415	-	-	-
60-105	138-228	139-355	213-394	-	-	-
PARSEC-II						
45-120	107-229	112-239	92-221	111-294	133-366	-
55-110	117-150 153-211	130-227	109-202	138-270	166-335	-
60-105	125-145 158-203	140-221	118-193	151-258	182-320	-
FRANEC						
45-120	111-262	113-272	136-415	183-565	220-600	-
55-110	131-242	134-251	173-378	233-514	282-600	-
60-105	141-232	145-240	192-360	259-488	313-592	-

$$\alpha_{GSMF} = -1.45 \longrightarrow \alpha_{GSMF}(z)$$



PISN / CCSN ratio


$$\frac{dN_{CCSN}}{dM_{SFR}} = \int_{8 M_{\odot}}^{50 M_{\odot}} \phi(M) dM$$

σ_Z	PI/CC ($z = 0$)	PI/CC ($z = z_{\text{peak}}^{\text{PI}}$)	PI/CC ($z = 6$)
variation P ($Z_{\text{max}} = 1.5 \times 10^{-3}$)			
0.15	2.5×10^{-9}	1.4×10^{-7}	2.3×10^{-7}
0.35	9.2×10^{-6}	3.5×10^{-5}	5.5×10^{-5}
0.70	1.7×10^{-4}	2.4×10^{-4}	3.3×10^{-4}
variation F ($Z_{\text{max}} = 6.6 \times 10^{-3}$)			
0.15	9.2×10^{-4}	2.3×10^{-3}	4.5×10^{-3}
0.35	2.2×10^{-3}	3.5×10^{-3}	5.2×10^{-3}
0.70	4.3×10^{-3}	5.4×10^{-3}	6.6×10^{-3}
variation O ($Z_{\text{max}} = 1.7 \times 10^{-2}$)			
0.15	1.5×10^{-2}	2.2×10^{-2}	2.8×10^{-2}
0.35	1.5×10^{-2}	2.0×10^{-2}	2.4×10^{-2}
0.70	1.5×10^{-2}	1.7×10^{-2}	1.9×10^{-2}

Host galaxy properties

$$M_{\star} \sim 10^8 - 10^9 M_{\odot}$$

$$\text{variation P} \\ \sigma_Z = 0.15$$




$$Z_{max} = 1.5 \times 10^{-3}$$

$$Z \lesssim Z_{max}$$

Host galaxy properties

$$M_{\star} \sim 10^{10} - 10^{11} M_{\odot}$$

$$\text{variation O} \\ \sigma_Z = 0.35$$



$$Z_{max} = 1.7 \times 10^{-2}$$

$$Z_{max} \sim Z_{peak}^{SFRD}$$



Pop III

peak of PISN rate at
 $Z \sim 10^{-2} - 10^{-3}$



Pop II stars main PISN progenitors, not Pop III

Pop III SFRD Hartwig et al. 2022





CAVEAT: SFRD and IMF at high z highly uncertain


$M_{up} = 300 M_{\odot}$ to $600 M_{\odot}$

$$\frac{dN_{PISN}}{dM_{SFR}}(Z)$$

PISN rate



FMR Mannucci et al. 2010



$$12 + \log(O/H) = 8.90 + 0.37m - 0.14s - 0.19m^2 + 0.12ms - 0.054s^2$$




$$m = \log(M_{\star}) - 10$$

$$s = \log(\text{SFR})$$




$$\log Z = 12 + \log(O/H) - 10.58$$

M_{He} criterion


$$\frac{dN_{PISN}}{dM_{SFR}}(Z)$$



FMR Mannucci et al. 2010



FMR Mannucci et al. 2011



$$12 + \log(\text{O}/\text{H}) = 8.90 + 0.37m - 0.14s - 0.19m^2$$

$$+ 0.12ms - 0.054s^2 \quad \text{for } \mu_{0.32} \geq 9.5$$

$$= 8.93 + 0.51(\mu_{0.32} - 10) \quad \text{for } \mu_{0.32} < 9.5,$$

$$\mu_{\alpha} = \log(M_{*}) - \alpha \log(\text{SFR})$$

$$m = \log(M_{*}) - 10$$

$$s = \log(\text{SFR})$$